# **Environmental** Science & lechnology

# Role of Anaerobic Ammonium Oxidation (Anammox) in Nitrogen Removal from a Freshwater Aquifer

Richard L. Smith,\*<sup>,†</sup> J. K. Böhlke,<sup>‡</sup> Bongkeun Song,<sup>§</sup> and Craig R. Tobias<sup>#</sup>

<sup>†</sup>National Research Program, U.S. Geological Survey, 3215 Marine Street, Boulder, Colorado 80303, United States

<sup>‡</sup>National Research Program, U.S. Geological Survey, 431 National Center, Reston, Virginia 20192, United States

<sup>§</sup>Virginia Institute of Marine Science, College of William and Mary, P.O. Box 1346, Gloucester Point, Virginia 23062, United States <sup>#</sup>Department of Marine Sciences, University of Connecticut, 1080 Shennecossett Road, Groton, Connecticut 06340, United States

**Supporting Information** 

**ABSTRACT:** Anaerobic ammonium oxidation (anammox) couples the oxidation of ammonium with the reduction of nitrite, producing N<sub>2</sub>. The presence and activity of anammox bacteria in groundwater were investigated at multiple locations in an aquifer variably affected by a large, wastewater-derived contaminant plume. Anammox bacteria were detected at all locations tested using 16S rRNA gene sequencing and quantification of hydrazine oxidoreductase (*hzo*) gene transcripts. Anammox and denitrification activities were quantified by *in situ* <sup>15</sup>NO<sub>2</sub><sup>-</sup> tracer tests along anoxic flow paths in areas of varying ammonium, nitrate, and organic carbon abundances. Rates of denitrification and anammox were determined by quantifying changes in <sup>28</sup>N<sub>2</sub>, <sup>29</sup>N<sub>2</sub>, <sup>30</sup>N<sub>2</sub>, <sup>15</sup>NO<sub>2</sub><sup>-</sup>, <sup>15</sup>NO<sub>2</sub><sup>-</sup>, and <sup>15</sup>NH<sub>4</sub><sup>+</sup> with groundwater travel time. Anammox was present and active in all areas tested, including where ammonium and dissolved organic carbon concentrations were low, but decreased in proportion to denitrification when acetate was added to increase available electron supply. Anammox contributed 39–90% of potential N<sub>2</sub> production in this aquifer,



with rates on the order of 10 nmol  $N_2$ -N  $L^{-1} day^{-1}$ . Although rates of both anammox and denitrification during the tracer tests were low, they were sufficient to reduce inorganic nitrogen concentrations substantially during the overall groundwater residence times in the aquifer. These results demonstrate that anammox activity in groundwater can rival that of denitrification and may need to be considered when assessing nitrogen mass transport and permanent loss of fixed nitrogen in aquifers.

## ■ INTRODUCTION

Humans have altered the global nitrogen (N) budget,<sup>1,2</sup> with impacts in virtually all aquatic and terrestrial environments. This includes freshwater aquifers, which receive an estimated 15 Tg/year or ~10% of all anthropogenic fixed N.<sup>2</sup> Denitrification is usually considered the key dissolved inorganic nitrogen (DIN) removal mechanism in groundwater,<sup>2,3</sup> but the discovery of anaerobic ammonium oxidation (anammox)<sup>4</sup> suggests that biogeochemical processes affecting the fate and transport of groundwater DIN, and thus its ultimate delivery to surface waters, may be far more complex than was previously thought. Relative rates of anammox and denitrification in groundwater are unknown, but could have far-reaching implications for N storage and water quality at local to global scales.

DIN (nitrate  $(NO_3^{-})$ , nitrite  $(NO_2^{-})$ , and ammonium  $(NH_4^{+})$ ) arrive in groundwater from atmospheric deposition and weathering of rocks and soil organic matter and from a variety of local anthropogenic sources including landfill leachate, wastewater disposal, septic systems, and agriculture. N fate and transport in shallow aquifers depend on local geology, recharge conditions, hydraulic conductivity, and inaquifer removal reactions. The protracted residence time of N

in aquifers, relative to surface freshwater, causes aquifers to have both large inventories of anthropogenic N and extended contact times for in-aquifer reactions to potentially attenuate N loads during transport.  $NO_3^-$  is typically the dominant form of DIN in oxic aquifers, moving freely through the unsaturated and saturated zone. NH4<sup>+</sup> tends to dominate the DIN pool in suboxic aquifers, but its transport is slower due to chemical sorption and exchange with aquifer sediments.<sup>5</sup> Coexistence of  $NO_3^-$  and  $NH_4^+$ , as well as  $O_2$  and organic-C, can occur in groundwater due to differential advection rates, resulting in mixed redox zones of dissimilatory N-cycling processes. Such zones may be characterized by co-occurrence of N species of multiple redox states (e.g.,  $NO_3^-$  and  $NH_4^+$ ) and production/ consumption of NO<sub>2</sub><sup>-</sup> through both oxidizing and reducing reactions.<sup>6</sup> Denitrification has long been considered the primary N removal mechanism of fixed N and has been examined in a number of subsurface studies,  $7^{-11}$  with an estimated 20% of the global extent of the process occurring in groundwater.<sup>3</sup>

Received:May 20, 2015Revised:September 16, 2015Accepted:September 24, 2015Published:September 24, 2015



**Figure 1.** Map of study site on Cape Cod, MA, USA, showing groundwater contaminant plume, selected well locations, tracer test site locations, and depth profiles at the tracer test locations of groundwater concentrations of  $NO_3^-$ ,  $NH_4^+$ , and  $O_2$  in August 2011 (B) and June 2012 (C). Depths and primary constituents for each tracer test are indicated in yellow boxes. Elevation is relative to mean sea level.

Typically, excess  $N_2$  production in aquifers is attributed exclusively to denitrification,<sup>10,12</sup> which is commonly organic-C driven, but can also utilize inorganic electron supplies, such as reduced iron and sulfur.<sup>11,13,14</sup>

Anammox couples NH4<sup>+</sup> oxidation with NO2<sup>-</sup> reduction to produce nitrogen gas  $(N_2)$  under suboxic to anoxic conditions. Provided there is a source of NO<sub>2</sub><sup>-</sup>, anammox can potentially remove large amounts of fixed N without requiring substantial amounts of organic-C, reduced iron, or mineral phase electron donors. Occurrence of anammox in environmental systems can have profound implications for DIN mass balance and natural attenuation. Anammox was first discovered in a wastewater treatment plant.<sup>4</sup> Since then, it has been found to be variably important in soils, in marine and fresh surface waters, and near sediment-water interfaces.<sup>15-22</sup> Its presence in groundwater has been inferred locally in contaminated settings from distributions of DIN constituents, natural abundance <sup>15</sup>N distributions, and biomarkers,<sup>23-25</sup> but has not been directly measured. In groundwater, anammox is not necessarily limited to narrow diffusional zones like it is in many surface water or soil environments; groundwater hydrology could create expanded zones suitable for anammox communities and widespread spatial distribution of anammox activity. Although the impact of anammox on the fate of DIN in aquifers could be substantial, there has not been a comprehensive study conducted to measure the rate and significance of anammox in the subsurface environment. Thus, we conducted in situ<sup>15</sup>N tracer experiments and molecular characterization of groundwater communities in a N-contaminated sand and gravel aquifer.

#### MATERIALS AND METHODS

**Study Site.** This study was conducted in different parts of an aquifer that was variably affected by a groundwater contaminant plume located on Cape Cod, MA, USA. The plume was created by land disposal of treated wastewater from ~1930 to 1995 onto a sand and gravel glacial outwash plain. It is contained within the unconfined, water table aquifer and measures >8 km long in the direction of groundwater flow, ~1 km wide, and ~30 m thick on a vertical axis. The plume is characterized by concentration gradients of organic and inorganic constituents that are typical of domestic wastewater, redox gradients, gradients of secondary, naturally occurring constituents, such as Fe and As, that have been mobilized by the altered geochemical environment, and gradients in microbial community composition. At essentially all locations, uncontaminated groundwater lies above and below the contaminant plume. The plume site has been intensively instrumented by the U.S. Geological Survey to conduct longterm investigations on the fate and transport of contaminants in an unconsolidated aquifer.<sup>26</sup>

Groundwater Sample Collection and Analyses. Well construction and sampling procedures are described elsewhere.<sup>27</sup> Groundwater was collected from multilevel sampling wells (MLSs) using a peristaltic pump. Specific conductance, pH, dissolved oxygen, and temperature were determined in the field immediately upon collection. N2O samples were collected with a syringe and injected through a needle into stoppered, He-flushed serum bottles containing NaOH as a preservative. Samples for N<sub>2</sub> analysis were pumped into 150 mL serum bottles, which were filled to overflowing for several bottle volumes at a slow pumping rate. Then, a pellet of KOH was added, and the bottles were stoppered under water using a syringe needle as vent for the displaced water and kept chilled at 4 °C.7 All other samples for geochemical analysis were filtered with 0.45  $\mu$ m capsule filters and preserved. Samples for nitrate, nitrite, acetate, and other anions were frozen. Samples for major cations (including ammonium) and dissolved total iron were acidified to pH <2 with concentrated  $H_2SO_4$  or HNO<sub>3</sub>, respectively.

Dissolved oxygen was determined colorimetrically and with a probe.<sup>7</sup> Anions, cations, and acetate were analyzed by ion chromatography.<sup>28</sup> N<sub>2</sub>O was analyzed by gas chromatography.<sup>29</sup> Dissolved total iron was analyzed by inductively coupled plasma atomic emission spectroscopy.<sup>30</sup> Dissolved organic carbon (DOC) was analyzed with a C analyzer following acidification and persulfate oxidation.<sup>31</sup>

**Tracer Tests.** *In situ* tracer tests were conducted at sites F575 and F593 (Figure 1), which were instrumented with an array of MLSs located in rows down-gradient from an injection MLS (Figure S1). The injection tests were similar to previously described tests with  $NH_4^{+,5}$  Briefly, a 200 L gas-impermeable

bladder was filled with ultrahigh-purity He and evacuated seven times to remove residual oxygen. Then, ~1 L of boiled, Hesparged DI water containing dissolved tracer salts [NaBr,  $Na^{15}NO_2$  (98+ atom %), and with or without  $NaC_2O_2H_3$  was pumped into the bladder, followed by 100 L of groundwater collected from the appropriate injection well port. Target final concentrations of Br<sup>-</sup>,  $NO_2^{-}$ , and acetate were ~1100, 200, and 200  $\mu$ M, respectively. During the filling process, the bladder was frequently agitated to ensure thorough mixing and was kept in a wading pool of water that was maintained at groundwater temperature using ice. After mixing, any gas remaining within the bladder was vented and the tracer solution pumped into the injection port, with two to three samples collected during the injection process. Caution was taken during the entire tracer preparation and injection process to avoid any contact of the solution with air. All tracer tests were conducted in anoxic zones ( $<5 \mu M O_2$ ) within the aquifer (Tables S1 and S2).

Groundwater samples were collected from the injection port and appropriate ports in the tracer MLS array with time as detailed above. Arrival of the tracer cloud at down-gradient MLS ports was determined in the field by monitoring specific conductance.

Detection and Phylogenetic Analysis of Anammox Bacteria. Groundwater sampling was conducted at four wells (S469, F575, F168, and F442) within the groundwater contamination plume (Figure 1). Water samples (4 L) from each well were filtered through 0.22  $\mu$ m Millipore Sterivex filters (Millipore Corp., Billerica, MA, USA) to concentrate microbial biomass using a peristaltic pump and immediately frozen. Genomic DNA was extracted from the filters using a Gentra Puregene Genomic DNA Purification Kit (Gentra Systems, Inc., Minneapolis, MN, USA) as described by Oates and Song.<sup>32</sup> Anammox bacterial 16S rRNA genes in the extracted DNA were amplified, cloned, and sequenced as described in the Supporting Information.

Quantification of hzo and nosZ Transcripts. The gene transcripts in anammox (hydrazine oxidoreductase, hzo) and denitrification (nitrous oxide reductase, nosZ) were analyzed in conjunction with the tracer tests. For tests 1 and 2, groundwater samples (1 L) were pumped through 0.22  $\mu$ m Sterivex filters, whereas the samples from tracer tests 3 and 4 were filtered through 5  $\mu$ m membranes to remove sediment particles before the cells were concentrated on 0.22  $\mu$ m Supor filters (Pall Life Sciences). All filters were immediately frozen in liquid nitrogen. Total RNA was extracted using a mirVAN miRNA isolation kit (Ambion) and treated with a TURBO DNA-free kit (Ambion) to remove genomic DNA. The treated RNA was used for cDNA synthesis using an iScript cDNA synthesis kit (Bio-Rad). Quantitative PCR of hzo and nosZ transcripts was conducted as described by Lisa et al.,<sup>19</sup> whereas 16S rRNA was quantified using the primers 341F and 685R. The ratio of hzo or nosZ transcript copies to 16S rRNA copies was calculated and used to compare different levels of gene expression in each tracer experiment.

**Isotopic Analyses.** Isotopic analyses of  $NO_3^-$  and  $NO_2^-$  were done by modified versions of the bacterial denitrifier method, in which compounds of interest are converted to  $N_2O$  for isotope-ratio mass spectrometry (IRMS). *Pseudomonas aureofaciens* was used to convert  $NO_3^- + NO_2^-$  to  $N_2O_3^{33}$  and *Stenotrophomonas nitritireducens* was used to convert  $NO_2^-$  to  $N_2O_3^{34}$  Calibrations were done by analyzing laboratory standards as samples with <sup>15</sup>N mole fractions ( $x^{15}N$ ) ranging from 0.0037 to 0.982.  $NH_4^+$  was separated from groundwater

by diffusion into NaHSO<sub>4</sub> traps and analyzed for  $\delta^{15}$ N after reaction with O<sub>2</sub> to produce N<sub>2</sub> for continuous-flow IRMS.<sup>35</sup> Isotopologues of dissolved N<sub>2</sub> (<sup>14</sup>N<sup>14</sup>N, <sup>14</sup>N<sup>15</sup>N, and <sup>15</sup>N<sup>15</sup>N) in groundwater were analyzed by multiple techniques. First, sample headspace gas was reacted with CuO + Cu<sub>2</sub>O and CaO in sealed tubes to remove O<sub>2</sub>, H<sub>2</sub>O, and CO<sub>2</sub> from N<sub>2</sub> for dual-inlet IRMS.<sup>5</sup> Alternatively, headspace gas was passed unreacted through a mole-sieve gas chromatograph, and the N<sub>2</sub> peak was analyzed in continuous-flow mode.<sup>36</sup> For both methods, IRMS responses at *m*/*z* 28, 29, and 30 were calibrated against air and air-saturated water samples and adjusted for NO artifacts at *m*/*z* 30.

For each sample, measured N<sub>2</sub> isotopologue ratios were converted to estimated concentrations of background N<sub>2</sub>, N<sub>2</sub> produced by anammox during the experiment, and N<sub>2</sub> produced by denitrification during the experiment, using equations similar to those derived previously and presented as a "comprehensive approach" for three-component isotopologue attribution.<sup>37</sup> Application of those equations included the following modifications and assumptions: (1) Measured isotopologue ratios of in situ background N2 (rather than atmospheric  $N_2$ ) were used as the nontracer baseline values. (2) Although unlabeled NO<sub>3</sub><sup>-</sup> at F593 (tests 3 and 4) had  $\delta^{15}$ N of +15 to 19%, the  $NO_3^-$  was not considered to contribute unlabeled N to N2 production directly, but only indirectly through isotope dilution of the  ${}^{15}$ N-labeled NO<sub>2</sub><sup>-</sup> pool, which was measured in each sample and included in the N2 production equations. (3)  $NO_2^-$  contributing to  $N_2$  produced by both anammox and denitrification in each sample during the tracer experiment was assumed to have had a time-integrated  $x^{15}N$  equal to the midpoint of the measured  $x^{15}NO_2^{-}$  value at time zero and the measured value at the sample time (important mainly for test 4, in which  $x^{15}NO_2^{-1}$  decreased substantially). Measured variations and uncertainties of  $\delta^{15}$ N values of NH<sub>4</sub><sup>+</sup> were incorporated in the equations, although they had negligible effects on the calculated contributions of anammox and denitrification to N2. Reported values of anammox N<sub>2</sub> outnumber reported values of denitrification N<sub>2</sub> because some dual-inlet N2 isotopic analyses were affected by  $N_2O$  that was converted to  $N_2$  by reaction with CuO + Cu<sub>2</sub>O during off-line sample preparation for dual-inlet IRMS. For tests 1, 2, and 3, detection limits for accumulated N<sub>2</sub> produced by anammox and denitrification were approximately 2-5 nmol  $L^{-1}$ , corresponding to accumulation rates of the order of approximately 0.5 nmol L<sup>-1</sup> day<sup>-1</sup> over the length of the experiments. For test 4, detection limits were similar, but estimated rates may have larger uncertainties, as  $x^{15}N$  of NO<sub>2</sub><sup>-</sup> was changing rapidly, and possibly not linearly as assumed.

Note that for tracer test results denitrification and anammox are defined as the net reduction of NO<sub>2</sub><sup>-</sup> through N<sub>2</sub>O to produce N<sub>2</sub> (all N atoms in N<sub>2</sub> from NO<sub>2</sub><sup>-</sup>) and net anaerobic NH<sub>4</sub><sup>+</sup> oxidation coupled with NO<sub>2</sub><sup>-</sup> reduction to produce N<sub>2</sub> (one atom each from NH<sub>4</sub><sup>+</sup> and NO<sub>2</sub><sup>-</sup>), respectively. Production of <sup>30</sup>N<sub>2</sub> from <sup>15</sup>NO<sub>2</sub><sup>-</sup> could result from any combination of classical denitrification,<sup>38</sup> chemodenitrification,<sup>39</sup> or anammox "disguised" as denitrification <sup>40,41</sup> and thus may differ, *sensu stricto*, from the definition of anammox and denitrification as interpreted from the phylogenetic analysis and gene transcript data.

#### RESULTS AND DISCUSSION

**Groundwater Geochemistry.** This study investigated groundwater anammox in a large (>8 km long) wastewater

Article



**Figure 2.** (Left) Phylogenetic tree of anammox bacterial 16S rRNA genes detected in groundwater samples collected from well locations S469, F575, F168, and F442. Bootstrap values are indicated at branch points. (Right) Comparison of transcriptional expression of hydrazine oxidoreductase (*hzo*) and nitrous oxide reductase (*nosZ*) genes as transcript copies normalized to total 16S rRNA copies in groundwater communities collected during the tracer tests (1–4). Error bars are  $\pm$  standard deviation of replicate samples.

contaminant plume located in an unconfined sand and gravel aquifer used for drinking water (Figure 1A). The plume was characterized by vertical and longitudinal concentration gradients of dissolved and sorbed constituents, including  $NO_3^-$ ,  $NH_4^+$ , dissolved  $O_2$ , and DOC. Typical vertical profiles of groundwater chemistry for the upper and middle regions of the plume are shown in Figure 1B,C. The long axis of the plume center can be divided into three regions. The upgradient third, closest to the source, had anoxic zones of  $NO_3^-$ , Fe(III), and Mn(IV) reduction, and  $NO_3^-$  and  $NH_4^+$  did not coexist in the same water. The middle third of the plume had anoxic water that contained both  $NO_3^-$  and  $NH_4^+$ . The downgradient third contained  $O_2$  and  $NO_3^-$  but little or no  $NH_4^+$ , which was less mobile because of ion exchange in the aquifer.

Sixty years of wastewater discharge contributed an estimated organic carbon load to the sand and gravel aquifer of 600 t.<sup>31</sup> A portion of that carbon was sorbed onto aquifer solids,<sup>42,43</sup> and a portion remained as DOC, which was subsequently transported down-gradient. DOC decreased in concentration during transport by continued sorption, degradation, and dispersion.<sup>42,44,45</sup> After 3 km of transport in the vicinity of well site F593 (see map Figure 1), DOC concentrations had decreased by approximately 80% and were primarily composed of recalcitrant compounds.<sup>45–47</sup> The number of free-living bacteria also decreased within 3 km of the source by about 10-fold,<sup>45</sup> reflecting the limited supply of reactive electron donors in the existing geochemical conditions present at F593, the well site for tracer tests 3 and 4.

In 1995, wastewater disposal was discontinued and the contaminant plume was allowed to dissipate by natural attenuation. Since that time, the concentrations and depthintegrated masses of  $NO_3^-$ ,  $NH_4^+$ , and DOC have substantially decreased within the 0.5 km interval down-gradient of the wastewater disposal beds,<sup>5,48</sup> which includes F575, the well site for tracer tests 1 and 2. In contrast, the changes in concentrations at F593 during this same period have been relatively small, particularly for ammonium (Figure S3), as have changes in the integrated mass of DOC.<sup>48</sup> Available electron supply from degradation of reactive organic compounds continued to be limiting at the time of the tracer tests at F593 and was in the process of decreasing at F575. Degradation and mineralization of organic compounds sorbed onto aquifer solids underneath the former disposal beds continued to consume  $O_2$  coming from up-gradient groundwater sources and were serving as a source of the continued supply of inorganic nitrogen arriving at F575.<sup>42</sup>

Microbial Evidence for Anammox and Denitrification. Anammox bacteria were detected by targeting 16S rRNA genes in groundwater obtained from the entire length of the NH<sub>4</sub><sup>+</sup>containing zone within the contaminant plume, which extended 3.3 km to well F271 in 2013 (Figure 1A). Anammox community composition was related to sampling location. Three different operational taxonomic units (OTUs) were present at the former wastewater disposal beds that were closely related to Candidatus Kuenenia and Brocadia spp. Kuenenia- and Brocadia-like sequences and have been reported in aquifers, wetlands, lakes, and terrestrial environments and likely represent anammox species that occur naturally in freshwater.<sup>20,22,25,49</sup> One of the anammox OTUs (S469 rep3) was unique to the disposal bed location; one (S469 rep2) clustered with other anammox bacteria collected 0.3 km downgradient; and the third (S469 rep1) clustered with other anammox bacteria in the middle region of the contaminant plume at 2.5 and 2.7 km down-gradient (Figure 2A).

Transcriptional expression of the genes involved in anammox and denitrification was detected at both tracer locations during all four tracer experiments. A key anammox gene, *hzo*, was uniformly expressed at both locations, but was highest for test 3 (Figure 2B). This showed uncoupling of *hzo* gene expression and activities of anammox bacteria, as previously reported.<sup>50</sup> On the basis of 16S rRNA gene sequences, Brocadia-like bacteria were present during tests 1 and 2, whereas bacteria associated with "Ca. Kuenenia spp." were expressed during tests 3 and 4. The gene expression for the final step of the denitrification pathway, N<sub>2</sub>O reductase (*nosZ*), indicated that the reduction capacity for N<sub>2</sub>O was more prevalent at the down-gradient location (tests 3 and 4).

**Rates of Anammox and Denitrification from** *in Situ* **Tracer Tests.** Natural gradient tracer tests were conducted within the aquifer using <sup>15</sup>NO<sub>2</sub><sup>-</sup> and Br<sup>-</sup> as tracers to demonstrate *in situ* activity and to investigate the competition between anammox and denitrification for electron acceptor supply and their relative contribution to overall N<sub>2</sub> production. Tests were conducted at plume locations 0.3 and 2.5 km from



**Figure 3.** Breakthrough curves of natural gradient tracer tests 1 (A, B) and 2 (C, D) at the injection port and a down-gradient sampling port: added tracer constituent concentrations normalized to injection concentrations ( $C/C_0$ ); other reactant and product concentrations; accumulated N<sub>2</sub> production attributed to anammox and denitrification; and <sup>15</sup>N mole fraction ( $x^{15}$ N) of recovered NO<sub>2</sub><sup>-</sup>. Travel distances for individual curves are indicated by horizontal bars above the figure panels; dotted lines represent 0 m, and solid lines represent 6 m. See Figure 1 and Figure S1 for maps of tracer test wells and tracer cloud travel paths. Error bars are ± one standard deviation of replicate samples.

the wastewater source, representing groundwater travel times of the order of 2 and 20 years, respectively.<sup>51,52</sup> These locations differed in labile C supply and exhibited two different anammox OTUs (F575 rep and F168 rep; Figure 2A). At the up-gradient location (F575) where more reactive C was available, parts of the vertical profile containing NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> did not overlap and were separated by an anoxic, DIN-free zone (Figure 1B). We hypothesized that the DIN-free zone was due to coupled denitrification/anammox activity that reduced NO<sub>3</sub><sup>-</sup> to NO<sub>2</sub><sup>-</sup> and then produced N<sub>2</sub> from NO<sub>2</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>. NO<sub>2</sub><sup>-</sup> is naturally present in groundwater just above the DIN-free zone (6.2-8.7 m elevation; Table S1). At the down-gradient location (F593), where reactive C was less abundant in the water or aquifer sediment,  $NO_3^-$  and  $NH_4^+$  coexisted in the absence of  $O_2$ throughout the vertical profile below about -5 m elevation (Figure 1C).

Four tracer tests were conducted. Tests 1 and 2 at the upgradient location compared activity within the  $NH_4^+$  zone and DIN-free zone, respectively. Tests 3 and 4 were performed in a single zone at the down-gradient location and compared the absence (test 3) and presence (test 4) of added acetate as an additional electron supply. Groundwater tracer travel distances ranged from 0 to 11 m (Figure S1), and travel times ranged from 0 to 40 days.

Anammox activity was detected in all four tests as  $^{29}N_2$  $(\pm^{28}N_2 \text{ depending on } x^{15}NO_2^-; \text{ Figures 3 and 4})$ , including test 2 in a zone in which no detectable dissolved  $NH_4^+$  was present. Rates of anammox  $N_2$  production were similar for tests 1-3, ranging from 9.1 to 12.2 nmol N  $L^{-1}$  day<sup>-1</sup> (Table 1), and 50fold higher in test 4. Denitrification activity was detected in all four tests as  ${}^{30}N_2$  ( $\pm {}^{29}N_2$  depending on  $x^{15}NO_2^{-}$ ) and as nitrous oxide ( $N_2O$ ). It was lowest in test 3, where labile C was scarce, and was stimulated 400-600-fold by acetate in test 4. With the exception of tracer test 4, where the rate of  $NO_3^{-1}$ reduction to  $NO_2^-$  was high, the <sup>15</sup>N mole fraction in the  $NO_2^$ tracer generally remained high with time and travel distance (Figures 3 and 4), indicating little or no detectable  ${}^{14}NO_{2}$ production.  $\delta^{15}NH_4^+$  values in representative tracer breakthrough peak samples from tests 3 and 4 were higher than background values by no more than 1 and 10%, respectively, indicating the rate of NH4<sup>+</sup> production by dissimilatory nitrate reduction to ammonium (DNRA) was not more than approximately 2 and 1% of the total N2 production rate for tests 3 and 4. For test 1,  $\delta^{15}NH_4^+$  measurements were affected by cross-contamination and not useful for assessing DNRA, but nonetheless confirmed that <sup>15</sup>N-enriched NH<sub>4</sub><sup>+</sup> significant source of error in the calculation of N2 components. Thus, anammox disguised as denitrification<sup>40,41</sup> was not a significant source of  ${}^{30}N_2$ , at least in tests 1, 3 and 4. NH<sub>4</sub><sup>+</sup> isotope ratios



**Figure 4.** Breakthrough curves of natural gradient tracer tests 3 (A, B) and 4 (C, D) at the injection port and at down-gradient sampling ports: added tracer constituent concentrations normalized to injection concentrations ( $C/C_0$ ); other reactant and product concentrations; accumulated N<sub>2</sub> production attributed to anammox and denitrification; and <sup>15</sup>N mole fraction ( $x^{15}$ N) of recovered NO<sub>2</sub><sup>-</sup>. Travel distances for individual curves are indicated by horizontal bars above the figure panels; dotted lines represent 0 m, dashed lines represent 3.5 m, and solid lines represent 7 or 11 m. See Figure 1 and Figure S1 for maps of tracer test wells and tracer cloud travel paths. Error bars are ± one standard deviation of replicate samples.

Table 1. Gas Production Rates by Anammox and Denitrification from in Situ Groundwater Tracer Tests Using <sup>15</sup>NO<sub>2</sub><sup>-a</sup>

	nitrogen gas (nmol N L <sup>-1</sup> day <sup>-1</sup> )		nitrous oxide (nmol N L <sup>-1</sup> day <sup>-1</sup> )	anammox contribution (%)	
tracer test	anammox	denitrification	(net)	N <sub>2</sub> produced	$N_2 + N_2O$ produced
1	12.2 (8.9)	3.9 (1.2)	107 (30.8)	75.7	9.9
2	10.7 (3.8)	16.6 (4.3)	20.4 (5.9)	39.0	22.3
3	9.1 (1.3)	1.0 (0.3)	1.6 (1.1)	89.7	77.8
4 <sup>b</sup>	458 (98.7)	662 (233)	324 (266)	40.9	31.7

<sup>*a*</sup>Rates were calculated from the average accumulated concentrations at all time points within the tracer breakthrough curves with the longest travel distance (see Figures 3 and 4) assuming constant accumulation with time during transport; values in parentheses are  $\pm$  one standard deviation. Rates do not include compensation for dispersion. <sup>*b*</sup>Sodium acetate was included with other tracers in test 4.

could not be measured in test 2 samples, which had undetectable  $NH_4^+$  concentrations. At the up-gradient site,  $N_2$ production via denitrification slightly exceeded  $N_2$  production by anammox in the shallower test near the  $NO_3^-$  zone (test 2) but was less than  $N_2$  production by anammox in the deeper  $NH_4^+$  zone (test 1; Table 1). However,  $N_2O$  was also produced by denitrification in tests 1 and 2.  $N_2O$  production rates differed between the two zones but in both cases exceeded  $N_2$ production. If denitrification is considered as the sum of  $N_2$  and  $N_2O$  production, then the relative proportion of anammox activity was lower (Table 1). There were small amounts of dissolved Fe(II) in the tracer test 1 zone (Table S1); thus, it is possible that some portion of the  $N_2O$  production in that test was due to chemodenitrification.<sup>39</sup> The largest relative contribution of anammox to total  $N_2$  production was approximately 90% in test 3 at the down-gradient location where the  $NH_4^+$  concentration was high and labile organic-C was scarce.

Spatial/Temporal Controls of Anammox and Denitrification Rates. When  $NO_2^-$  was supplied as a tracer,  $N_2$ production by anammox was detectable even within a few hours at the injection wells. If relatively abundant degradable C was present (tests 1, 2, and 4), denitrification also was detected immediately. But when denitrification was limited by more restricted electron supply (test 3), 90% of the  $N_2$  production was attributable to anammox; this is higher than the relative anammox contribution reported in almost all other environments studied to date. The addition of an exogenous electron supply, in the form of acetate (test 4; Figure 4C,D), substantially stimulated both anammox and denitrification. Anammox bacteria can utilize short-chain organic acids to enhance growth.<sup>41</sup> In the presence of acetate, NO<sub>3</sub><sup>-</sup> was the primary oxidant and NO<sub>2</sub><sup>-</sup> production and consumption rates were nearly balanced, resulting in decreasing <sup>15</sup>NO<sub>2</sub><sup>-</sup> mole fraction due to <sup>14</sup>NO<sub>3</sub><sup>-</sup> reduction to <sup>14</sup>NO<sub>2</sub><sup>-</sup>; 100% of the NO<sub>3</sub><sup>-</sup> decrease was recovered as N<sub>2</sub>O and N<sub>2</sub>. The anammox stimulation in test 4 relative to test 3 indicated enhanced anaerobic NH<sub>4</sub><sup>+</sup> oxidation in the presence of the organic-*C*, in essence representing a multiplier effect for total N<sub>2</sub> production beyond what might be predicted stoichiometrically from oxidation of organic-C by denitrification alone.

The timing of the anammox and denitrification  $N_2$  product peaks relative to the arrival of the conservative tracer demonstrated interesting interactions between the two processes. During test 1 at the up-gradient location in the zone in which  $NH_4^+$  was present but N oxides were not and likely had not been for many years (Figure S2), the highest rates of  $NH_4^+$  oxidation to  $N_2$  were evident within the leading edge of the tracer cloud, whereas denitrification took longer to respond and the highest concentration of  $N_2$  from denitrification occurred at the end of the tracer breakthrough peak. This suggests that anaerobic  $NH_4^+$  oxidation was active in this zone and quickly responded to the  $NO_2^-$  tracer, whereas denitrifying activity required time for induction.

Tracer test 2 was conducted in a transient zone in which vertical gradients of  $NO_3^-$ ,  $NH_4^+$ , and DOC concentrations were changing at interannual and possibly shorter time scales (Figure S2). These changes were driven by a combination of fluctuations in the water table elevation (and associated fluctuations in gradient direction) and variations in up-gradient sources of groundwater constituents. This likely resulted in a subsurface microbial population capable of responding to the fluctuating N speciation. Anammox and denitrification occurred concurrently and consistently during test 2, at approximately equivalent rates, with relatively little  $N_2O$  net accumulation. This appears to be a zone primed for removing fixed N, with a low but steady production of electrons and  $NH_4^+$  via degradation of the organic pool.

In contrast, tests 3 and 4 were conducted in a C-depleted zone that contained high concentrations of both NO3- and  $NH_4^+$  (Figure S3 and Table S2). Anammox was clearly active and ready to respond rapidly to the tracers at this site. In test 3, N<sub>2</sub> accumulation was nearly linear with time, suggesting a zeroorder response to the NO<sub>2</sub><sup>-</sup> addition, even at relatively low concentrations (<10  $\mu$ M) in the trailing edge of the tracer cloud (Figure 4B). A previous tracer test in this zone with <sup>15</sup>NH<sub>4</sub><sup>+</sup> determined that anammox activity was below an estimated detection limit of 27 nmol N L<sup>-1</sup> day<sup>-1.5</sup> This is consistent with the anammox rate of 9.1 nmol N  $\rm \dot{L}^{-1}~day^{-1}$  in the current study using <sup>15</sup>NO<sub>2</sub><sup>-</sup>, which permitted quantification with an improved detection limit. Denitrification at this location was undetectable until an electron supply was added, after which both anammox and denitrification activity responded vigorously, in a short time interval (days) relative to the contaminant travel time to this site (decades).

**Implications for Anammox in Aquifers.** This study is one of only a few giving evidence for anammox potential in aquifers and the first to quantify its rate and relative importance with respect to denitrification. If anammox is found to be

common in freshwater aquifers, it could have a substantial effect on local, regional, and global assessments of DIN fluxes and the return of fixed N to the atmosphere. Anammox was clearly present and active in Cape Cod groundwater under a variety of geochemical conditions. The process was not necessarily limited to narrow zones by diffusive flux, but rather was found in situations where fixed N persisted, whether oxidized or reduced, and dissolved O2 was depleted. These are conditions commonly found in the subsurface and suggest that anammox can be an important geochemical mediator in that environment. This could be especially true where groundwater has been affected by wastewater,  $2^{23-25}$  whether from domestic, agricultural, or industrial sources, including hydrocarbon extraction, in heterogeneous aquifers where flow velocities and redox reaction rates vary locally and reaction zones overlap,<sup>53</sup> in groundwater recharge areas beneath wetlands or lakes,<sup>20,54,55</sup> and in groundwater discharge areas (riparian zones) where oxic and suboxic groundwater flow paths converge and encounter various lithologies.<sup>11,56</sup> This study indicates that although anammox contribution can range widely, depending on geochemistry, it can be a significant contributor to in situ DIN removal and total N<sub>2</sub> production in freshwater aquifers. Anammox appears to be favored when the ratio of NO3<sup>-</sup> to degradable DOC is high.<sup>57–59</sup> Thus, whereas subsurface rates of activity may be substantially lower than those at the surface water-land interface, groundwater residence times commonly are much longer than in surface freshwater, and the volume of affected water may be greater, resulting in potentially equivalent or even greater total N removal in the subsurface.

#### ASSOCIATED CONTENT

#### **S** Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.5b02488.

PCR amplification method, maps of the tracer test sites (Figure S1),  $NO_3^-$  and  $NH_4^+$  concentration histories at sites F575 (Figure S2) and F168 (Figure S3), and groundwater geochemistry at the time of the tracer tests at F575 (Table S1) and F168 (Table S2) (PDF)

#### AUTHOR INFORMATION

#### **Corresponding Author**

\*(R.L.S.) Phone: (303) 541-3032; fax: (303) 541-3084; e-mail: rlsmith@usgs.gov.

### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

This study was funded by National Science Foundation Grants EAR 1024662, 1329284, and 1024900, the USGS Toxics Substances Hydrology Program, and the USGS National Research Program. Field site logistics and coordination were provided by D. LeBlanc and T. McCobb. Assistance with field sample collection was provided by D. A. Repert, J. B. Taylor, C. T. Buckner, N. W. Chang, and V. Rollinson. Isotope analyses were done by S. Mroczkowski and J. Hannon; molecular characterization was done by J. B. Taylor, T. Sparrer, and A. Zoumplis; solute chemical analyses were done by D. A. Repert. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

#### REFERENCES

(1) Galloway, J. N.; Dentener, F. J.; Capone, D. G.; Boyer, E. W.; Howarth, R. W.; Seitzinger, S. P.; Asner, G. P.; Cleveland, C. C.; Green, P. A.; Holland, E. A.; Karl, D. M.; Michaels, A. F.; Porter, J. H.; Townsend, A. R.; Vörösmarty, C. J. Nitrogen cycles: Past, present, and future. Biogeochemistry 2004, 70, 153-226.

(2) Schlesinger, W. H. On the fate of anthropogenic nitrogen. Proc. Natl. Acad. Sci. U. S. A. 2009, 106, 203-208.

(3) Seitzinger, S.; Harrison, J. A.; Böhlke, J. K.; Bouwman, A. F.; Lowrance, R. R.; Peterson, B.; Tobias, C.; van Drecht, G. Denitrification across landscapes and waterscapes: A synthesis. Ecol. Appl. 2006, 16, 2064-2090.

(4) Mulder, A.; van de Graaf, A. A.; Robertson, L. A.; Kuenen, J. G. Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. FEMS Microbiol. Ecol. 1995, 16, 177-184.

(5) Böhlke, J. K.; Smith, R. L.; Miller, D. N. Ammonium transport and reaction in contaminated groundwater: Application of isotope tracers and isotope fractionation studies. Water Resour. Res. 2006, 42, W05411.

(6) Smith, R. L.; Yoshinari, T. Occurrence and turnover of nitric oxide in a nitrogen-impacted sand and gravel aquifer. Environ. Sci. Technol. 2008, 42, 8245-8251.

(7) Smith, R. L.; Böhlke, J. K.; Garabedian, S. P.; Revesz, K. M.; Yoshinari, T. Assessing denitrification in groundwater using natural gradient tracer tests with <sup>15</sup>N: In situ measurement of a sequential multistep reaction. Water Resour. Res. 2004, 40, W07101.

(8) Hiscock, K. M.; Lloyd, J. W.; Lerner, D. N. Review of natural and artificial denitrification of ground water. Water Res. 1991, 25, 1099-1111.

(9) Korom, S. F. Natural denitrification in the saturated zone: A review. Water Resour. Res. 1992, 28, 1657-1668.

(10) Vogel, J. C.; Talma, A. S.; Heaton, T. H. E. Gaseous nitrogen as evidence for denitrification in groundwater. J. Hydrol. 1981, 50, 191-200.

(11) Böhlke, J. K.; Wanty, R.; Tuttle, M.; Delin, G.; Landon, M. Denitrification in the recharge area and discharge area of a transient agricultural nitrate plume in a glacial outwash sand aquifer, Minnesota. Water Resour. Res. 2002, 38, 1105.

(12) Groffman, P. M.; Altabet, M. A.; Böhlke, J. K.; Butterbach-Bahl, K.; David, M. B.; Giblin, A. E.; Kana, T. M.; Nielsen, L. P.; Firestone, M. K.; Voytek, M. A. Methods for measuring denitrification: diverse approaches to a difficult problem. Ecol. Appl. 2006, 16, 2091-2122.

(13) Kölle, W.; Strebel, O.; Böttcher, J. Formation of sulfate by microbial denitrification in a reducing aquifer. Water Supply 1985, 3, 35 - 40.

(14) Postma, D.; Boesen, C.; Kristiansen, H.; Larsen, F. Nitrate reduction in an unconfined sandy aquifer: Water chemistry, reduction processes, and geochemical modeling. Water Resour. Res. 1991, 27, 2027-2045.

(15) Dalsgaard, T.; Canfield, D. E.; Petersen, J.; Thamdrup, B.; Acuña-González, J. N2 production by the anammox reaction in the anoxic water column of the Golf Dulce, Costa Rica. Nature 2003, 422, 606 - 608

(16) Thamdrup, B.; Dalsgaard, T.; Jensen, M. M.; Ulloa, O.; Farias, L.; Escribano, R. Anaerobic ammonium oxidation in the oxygendeficient waters off nothern Chile. Limnol. Oceanogr. 2006, 51, 2145-2156.

(17) Kuypers, M. M. M.; Sliekers, A. O.; Lavik, G.; Schmid, M.; Jorgensen, B. B.; Kuenen, J. G.; Damste, J. S. S.; Strous, M.; Jetten, M. S. M. Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. Nature 2003, 422, 608-611.

(18) Trimmer, M.; Nicholls, J. C.; Deflandre, B. Anaerobic ammonium oxidation measured in sediments along the Thames Estuary, United Kingdom. Appl. Environ. Microbiol. 2003, 69, 6447-6454.

(19) Lisa, J. A.; Song, B.; Tobias, C. R.; Duernberger, K. A. Impacts of freshwater flushing on anammox community structure and activities in the New River Estuary, USA. Aquat. Microb. Ecol. 2014, 72, 17-31. Article

den Camp, H. J. M.; Risgaard-Petersen, N.; Schwark, L.; Peng, Y.; Hefting, M. M.; Jetten, M. S. M.; Yin, C. Hotspots of anaerobic ammonium oxidation at land-freshwater interfaces. Nat. Geosci. 2013, 6, 103-107.

(21) Erler, D. V.; Eyre, B. D.; Davison, L. The contribution of anammox and denitrification to sediment N2 production in a surface flow constructed wetland. Environ. Sci. Technol. 2008, 42, 9144-9150.

(22) Wenk, C. B.; Blees, J.; Zopfi, J.; Veronesi, M.; Bourbonnais, A.; Schubert, C. J.; Niemann, H.; Lehmann, M. F. Anaerobic ammonium oxidation (anammox) bacteria and sulfide-dependent denitrifiers coexist in the water column of a meromictic south-alpine lake. Limnol. Oceanogr. 2013, 58, 1-12.

(23) Clark, I.; Timlin, R.; Bourbonnais, A.; Jones, K.; Lafleur, D.; Wickens, K. Origin and fate of industrial ammonium in anoxic ground water - <sup>15</sup>N evidence for anaerobic oxidation (anammox). Groundwater Monit. Rem. 2008, 28, 73-82.

(24) Robertson, W. D.; Moore, T. A.; Spoelstra, J.; Li, L.; Elgood, R. J.; Clark, I. D.; Schiff, S. L.; Aravena, R.; Neufeld, J. D. Natural attenuation of septic system nitrogen by anammox. Groundwater 2012, 50, 541-553.

(25) Moore, T. A.; Xing, Y.; Lazenby, B.; Lynch, M. D. J.; Schiff, S.: Robertson, W. D.; Timlin, R.; Lanza, S.; Ryan, M. C.; Aravena, R.; Fortin, D.; Clark, I. D.; Neufeld, J. D. Prevalence of anaerobic ammonium-oxidizing bacteria in contaminated groundwater. Environ. Sci. Technol. 2011, 45, 7217-7225.

(26) Cape Cod Toxic-Substances Hydrology Research Site, Toxic Substances Hydrology Program; http://ma.water.usgs.gov/ MMRCape.

(27) Savoie, J. G.; Smith, R. L.; Kent, D. B.; Hess, K. M.; LeBlanc, D. R.; Barber, L. B. Ground-water quality data for a treated wastewater plume undergoing natural restoration, Ashumet Valley, Cape Cod Massachusetts, 1994–2004. In U. S. Geological Survey Data Series 198; U.S. GPO: Washington, DC, USA, 2006; p 339

(28) Smith, R. L.; Baumgartner, L. K.; Miller, D. N.; Repert, D. A.; Böhlke, J. K. Assessment of nitrification potential in ground water using short term, single-well injection experiments. Microb. Ecol. 2006, 51, 22-35.

(29) Brooks, M. H.; Smith, R. L.; Macalady, D. L. Inhibition of existing denitrification enzyme activity by chloramphenicol. Appl. Environ. Microbiol. 1992, 58, 1746-1753.

(30) Savoie, J. G.; Kent, D. B.; Smith, R. L.; LeBlanc, D. R.; Hubble, D. W. Changes in groundwater quality near two granular-iron permeable reactive barriers in a sand and gravel aquifer, Cape Cod, Massachusetts, 1997-2000. U. S. Geological Survey Water-Resources Investigations Report 03-4309; U.S. GPO: Washington, DC, USA, 2004; 77 pp.

(31) Repert, D. A.; Barber, L. B.; Hess, K. M.; Keefe, S. H.; Kent, D. B.; LeBlanc, D. R.; Smith, R. L. Long-term natural attenuation of carbon and nitrogen within a groundwater plume after removal of the treated wastewater source. Environ. Sci. Technol. 2006, 40, 1154-1162.

(32) Oates, H.; Song, B. Molecular detection of dissimilatory arsenate-respiring bacteria in North Carolina groundwater. In Arsenic Contamination of Groundwater: Mechanism, Analysis, and Remediation; Ahuja, S., Ed.; Wiley: New York, 2008; pp 83-94.

(33) Sigman, D. M.; Casciotti, K. L.; Andreani, M.; Barford, C.; Galanter, M.; Böhlke, J. K. A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater. Anal. Chem. 2001, 73, 4145-4153

(34) Böhlke, J. K.; Smith, R. L.; Hannon, J. E. Isotopic analysis of N and O in nitrite and nitrate by sequential selective bacterial reduction to N2O. Anal. Chem. 2007, 79, 5888-5895.

(35) Holmes, R. M.; McClelland, J. W.; Sigman, D. M.; Fry, B.; Peterson, B. J. Measuring <sup>15</sup>NH<sub>4</sub><sup>+</sup> in marine, estuarine, and fresh waters: an adaptation of the ammonia diffusion method for samples with low ammonium concentrations. Mar. Chem. 1998, 60, 235-243. (36) Nielsen, L. P. Denitrification in sediment determined from nitrogen isotope pairing. FEMS Microbiol. Lett. 1992, 86, 357-362.

(37) Spott, O.; Stange, C. F. A new mathematical approach for calculating the contribution of anammox, denitrification and atmosphere to an  $N_2$  mixture based on a <sup>15</sup>N tracer technique. *Rapid Commun. Mass Spectrom.* **2007**, *21*, 2398–2406.

(38) Tiedje, J. M. Denitrification. In *Methods of Soil Analysis Part 2. Chemical and Microbiological Properties*; Page, A. L., Miller, R. H., Keeney, D. R., Eds.; American Society of Agronomy and Soil Science Society of America: Madison, WI, USA, 1982; pp 1011–1026.

(39) Jones, L. C.; Peters, B.; Pacheco, J. S. L.; Casciotti, K. L.; Fendorf, S. Stable isotopes and iron oxide mineral products as markers of chemodenitrification. *Environ. Sci. Technol.* **2015**, *49*, 3444–3452.

(40) Kartal, B.; Kuypers, M. M. M.; Lavik, G.; Schalk, J.; Op den Camp, H. J. M.; Jetten, M. S. M.; Strous, M. Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium. *Environ. Microbiol.* **2007**, *9*, 635–642.

(41) Kartal, B.; van Niftrik, L. A.; Keltjens, J. T.; Op den Camp, H. J. M.; Jetten, M. S. M. Anammox – Growth physiology, cell biology, and metabolism. *Adv. Microb. Physiol.* **2012**, *60*, 211–262.

(42) Smith, R. L.; Repert, D. A.; Barber, L. B.; LeBlanc, D. R. Longterm groundwater contamination after source removal – The role of sorbed carbon and nitrogen on the rate of reoxygenation of a treatedwastewater plume on Cape Cod, MA, USA. *Chem. Geol.* **2013**, 337– 338, 38–47.

(43) Barber, L. B., II; Thurman, E. M.; Runnells, D. D. Geochemical heterogeneity in a sand and gravel aquifer: Effect of sediment mineralogy and particle size on the sorption of chlorobenzenes. *J. Contam. Hydrol.* **1992**, *9*, 35–54.

(44) Barber, L. B., II. Hierarchical analytical approach to evaluating the transport and biogeochemical fate of organic compounds in sewage-contaminated groundwater, Cape Cod, Massachusetts. In *Groundwater Contamination and Analysis at Hazardous Waste Sites;* Lesage, S., Jackson, R. E., Eds.; Dekker: New York, 1992; pp 73–120.

(45) Harvey, R. W.; Barber, L. B., II Associations of free-living bacteria and dissolved organic compounds in a plume of contaminated groundwater. *J. Contam. Hydrol.* **1992**, *9*, 91–103.

(46) Thurman, E. M.; Barber, L. B., II; LeBlanc, D. R. Movement and fate of detergents in groundwater: A field study. *J. Contam. Hydrol.* **1986**, *1*, 143–161.

(47) LeBlanc, D. R. Sewage plume in a sand and gravel aquifer, Cape Cod, Massachusetts. In *U. S. Geological Survey Water Supply Paper* 2218; U.S. GPO: Washington, DC, USA, 1984; p 28.

(48) Miller, D. N.; Smith, R. L. Microbial characterization of nitrification in a shallow, nitrogen-contaminated aquifer, Cape Cod, Massachusetts and detection of a novel cluster associated with nitrifying Betaproteobacteria. *J. Contam. Hydrol.* **2009**, *103*, 182–193.

(49) Humbert, S.; Tarnawski, S.; Fromin, N.; Mallet, M.-P.; Aragno, M.; Zopfi, J. Molecular detection of anammox bacteria in terrestrial ecosystems: distribution and diversity. *ISME J.* **2010**, *4*, 450–454.

(50) Park, H.; Rosenthal, A.; Ramalingam, K.; Fillos, J.; Chandran, K. Linking community profiles, gene expression and N-removal in anammox bioreactors treating municipal anaerobic digestion reject water. *Environ. Sci. Technol.* **2010**, *44*, 6110–6116.

(51) LeBlanc, D. R.; Garabedian, S. P.; Hess, K. M.; Gelhar, L. W.; Quadri, R. D.; Stollenwerk, K. G.; Wood, W. W. Large-scale natural gradient tracer test in sand and gravel, Cape Cod, Massachusetts: 1. Experimental design and observed tracer movement. *Water Resour. Res.* **1991**, *27*, 895–910.

(52) Shapiro, S. D.; LeBlanc, D. R.; Schlosser, P.; Ludin, A. Characterizing a sewage plume using the  ${}^{3}$ H- ${}^{3}$ He dating technique. *Groundwater* **1999**, *37*, 861–878.

(53) Green, C. T.; Böhlke, J. K.; Bekins, B. A.; Phillips, S. Mixing effects on apparent reaction rates and isotope fractionation during denitrification in a heterogenous aquifer. *Water Resour. Res.* **2010**, *46*, W08525.

(54) Katz, B. G.; Plummer, L. N.; Busenberg, E.; Revesz, K. M.; Jones, B. F.; Lee, T. M. Chemical evolution of groundwater near a sinkhole lake, Northern Florida: 2. Chemical patterns, mass transfer modeling, and rates of mass transfer reactions. *Water Resour. Res.* **1995**, *31*, 1565–1584. (55) Denver, J. M.; Ator, S. W.; Lang, M. W.; Fisher, T. R.; Gustafson, A. B.; Fox, R.; Clune, J. W.; McCarty, G. W. Nitrate fate and transport through current and former depressional wetlands in an agricultural landscape, Choptank Watershed, Maryland, United States. *J. Soil Water Conserv.* **2014**, *69*, 1–16.

(56) Vidon, P.; Hill, A. R. Denitrification and patterns of electron donors and acceptors in eight riparian zones with contrasting hydrogeology. *Biogeochemistry* **2004**, *71*, 259–283.

(57) Kraft, B.; Tegetmeyer, H. E.; Sharma, R.; Klotz, M. G.; Ferdelman, T. G.; Hettich, R. L.; Geelhoed, J. S.; Strous, M. The environmental controls that govern the end product of bacterial nitrate respiration. *Science* **2014**, *345*, 676–679.

(58) Babbin, A. R.; Ward, B. B. Controls on nitrogen loss processes in Chesapeake Bay sediments. *Environ. Sci. Technol.* **2013**, 47, 4189– 4196.

(59) Algar, C. K.; Vallino, J. J. Predicting microbial nitrate reduction pathways in coastal sediments. *Aquat. Microb. Ecol.* **2014**, *71*, 223–238.